

# Chicken Withdrawal Time Chart

This chart uses data from the **Compendium of Veterinary Products** database, March 30, 2007. We recommend that any user of a product refers to the label.

The following pages contain a reference table listing product names alphabetically. This document summarizes information on withdrawal times for food products (meat, eggs) according to the species and the route of administration.

The withdrawal times listed correspond to label dosages and directions. Deviations from label recommendations may lead to drug residues in food products.

When a label clearly states that the product requires a “zero withdrawal time” or that “no withdrawal time is necessary”, the product will be included in the chart and “**Od**” will be indicated. However, if a product label makes no mention of a withdrawal period, then this product will be excluded.

**Withdrawal Notes** are included if a text explanation is also needed to clarify the withdrawal statement. Egg withdrawal information can be found here.

Further information regarding the products listed here may be found in the product labels. If in doubt after reading the label, contact the manufacturer. Addresses and telephone numbers are listed in our Manufacturer’s A-Z index.

Every effort has been made to ensure the accuracy of the information presented. However, it remains the responsibility of the readers to familiarize themselves with the information contained on the product label or package insert. The publisher, editorial team, and all those involved in the production of this website (or CD) cannot be held responsible for publication errors or any consequence that could result from the use of presented information.

## Withdrawal Time Chart Abbreviations

aer/sp	aerosol spray	I.M.M.	intramammary	I.V.	intravenous
bd	beak dip	I.N.	intranasal	mr	milk replacer
dw	drinking water	I.O.	intra-ocular	S.C.	subcutaneous
Epdl.	epidural	I.P.	intraperitoneal	ts	thigh-stab
lmsn.	by immersion	I.R.	intraruminal	ww	wing web
Inhl.	by inhalation	I.Syn.	intrasynovial	d	days
lfltn.	by infiltration	I.Ten.	intratendinous	h	hours
I.A.	intra-articular	I.Tum.	intratumoral	w	weeks
I.Derm.	intra-dermal	I.U.	intra-uterine		
I.M.	intramuscular	I.Vag.	intravaginal		

## Chicken Withdrawal Time Chart

Withdrawal times listed correspond to label dosages only.

Product Name	Company Name	Species	Route of Administration	Meat	Milk
3-Nitro® 20%	Alpharma	Chickens	In the feed	5d	
89/03®	Intervet	Chickens	In ovo, S.C.	21d	
AE-Pox™ Vac	Wyeth Animal Health	Chickens	ww	21d	
AE-Vac™	Wyeth Animal Health	Chickens	dw	21d	
Amprol® 25% Feed Mix	Merial	Chickens	In the feed		
<b>Note:</b> Do not feed to laying birds in production.					
Aureomycin® 110 G	Alpharma	Chickens	In the feed	7d	
Aureomycin® 220 G	Alpharma	Chickens	In the feed	7d	
Ava-Bron® HN-63	Schering-Plough	Chickens	dw, I.O.	21d	
Avatec® Medicated Premix	Alpharma	Chickens	In the feed		
<b>Note:</b> Do not feed to replacement, laying or breeding chickens.					
Bio-Cox® 120 G	Alpharma	Chickens	In the feed		
<b>Note:</b> Do not feed to replacement, laying or breeding chickens.					
Bio-Cox® 120 G Plus	Alpharma	Chickens	In the feed		
<b>Note:</b> Do not feed to replacement, laying or breeding chickens.					
Biosol® Liquid	Pfizer	Chickens	dw	7-14d	
<b>Note:</b> Withdraw 7 days before slaughter of broilers and 14 days before slaughter of laying hens.					
Bio-Trach™	Intervet	Chickens	aer/sp, dw, I.O.	21d	
Booster P S Conc.	Jaapharm	Chickens	dw	5d	
<b>Note:</b> Eggs obtained from treated birds within 5 days after the latest treatment with this drug must not be used in food.					
Breedervac-III-Plus®	Intervet	Chickens	I.M., S.C.	42d	
Breedervac-IV-Plus®	Intervet	Chickens	I.M., S.C.	42d	
Breedervac-Reo-Plus®	Intervet	Chickens	I.M., S.C.	42d	
Broilerbron® B <sub>1</sub>	Schering-Plough	Chickens	aer/sp, I.O.	21d	
Bronchitis Vaccine (Mass. & Conn. Types)	Wyeth Animal Health	Chickens	aer/sp, dw	21d	
Bronchitis Vaccine (Mass. Type)	Wyeth Animal Health	Chickens	aer/sp, dw	21d	
Bronchitis Vaccine (Mass. Type) (Freeze-Dried)	Merial	Chickens	aer/sp, dw, I.O.	21d	
Bronchitis Vaccine (Mass. Type) (Frozen)	Merial	Chickens	aer/sp	21d	
Bron-Newcavac-M®	Intervet	Chickens	I.M., S.C.	42d	
Bursa Blen® M	Merial	Chickens	dw	21d	
Bursa Guard N-B-R	Merial	Chickens	I.M., S.C.	42d	
Bursa Guard Reo	Merial	Chickens	I.M., S.C.	42d	
Bursal Disease-Marek's Disease Vaccine (Standard, Chicken and Turkey)	Merial	Chickens	In ovo, S.C.	21d	
Bursal Disease-Marek's Disease Vaccine (Standard, Turkey)	Merial	Chickens	In ovo, S.C.	21d	
Bursal Disease Vaccine (S-706)	Merial	Chickens	aer/sp, dw, S.C.	21d	
Bursa-Vac®	Schering-Plough	Chickens	dw	21d	
Bursa-Vac® 3	Schering-Plough	Chickens	dw, I.O.	21d	
Bursine®-2	Wyeth Animal Health	Chickens	dw	21d	
Cav-Vac®	Intervet	Chickens	ww	21d	
<b>Note:</b> Do not vaccinate chickens in lay.					
Chlorosol-50	Vétoquinol	Chickens	In the feed	7d	
Cholervac-PM-1™	Intervet	Chickens	ww	21d	
<b>Note:</b> Do not vaccinate 35 days prior to onset of lay or during egg production.					
Clonevac D-78®	Intervet	Chickens	aer/sp, dw	21d	
Coban® Premix	Elanco	Chickens	In the feed		
<b>Note:</b> Do not feed to replacement and laying chickens.					
Coccivac®-B	Schering-Plough	Chickens	aer/sp, dw, In the feed	21d	
Coccivac®-D	Schering-Plough	Chickens	aer/sp, dw, In the feed	21d	
Combovac-30™	Intervet	Chickens	aer/sp, dw, I.O.	21d	
Co-op® Poly-Tonine A Super® Booster No. 1	IPCO	Chickens	dw	5d	
<b>Note:</b> Eggs obtained from treated birds within 5 days after the latest treatment with this drug must not be used in food.					
Coxistac® 6% Premix	Phibro	Chickens	In the feed		
<b>Note:</b> Do not feed to laying chickens.					
Coxistac® 12% Premix	Phibro	Chickens	In the feed		
<b>Note:</b> Do not feed to laying chickens.					
Coyden® 25	Huvepharma AD	Chickens	In the feed		
<b>Note:</b> Do not feed clopidol to caged layer replacements after 16 weeks of age. Do not give to laying hens.					
Cygro® 1% Premix	Alpharma	Chickens	In the feed	5d	
<b>Note:</b> Do not feed to replacement, laying or breeding chickens.					
Deccox® 6% Premix	Alpharma	Chickens	In the feed		
<b>Note:</b> Do not feed to laying birds in production.					
Dusting Powder	Dominion	Chickens	Premise, Topical	7d	
Ectiban® 25 Fly Killer	Engage	Chickens	Topical	7d	

**Withdrawal times listed correspond to label dosages only.**

**Chicken Withdrawal Time Chart**

Product Name	Company Name	Species	Route of Administration	Meat	Milk
Enterovax®	Schering-Plough	Chickens	aer/sp, dw	21d	
<i>Note:</i> Do not administer this product to breeders 18 weeks of age or older.					
Flavomycin® 4	Huvepharma AD	Chickens	In the feed		
<i>Note:</i> Do not feed to replacement, laying or breeding chickens.					
FP-Vac™	Intervet	Chickens	ww	21d	
<i>Note:</i> Do not vaccinate chickens within 4 weeks of onset of egg production or during egg production.					
Gallimune™ NC-BR	Merial	Chickens	I.M., S.C.	42d	
Gallimycin®	Vétoquinol	Chickens	dw	24h	
<i>Note:</i> Do not use in birds producing eggs for food purposes.					
Gallimycin®-50	Vétoquinol	Chickens	In the feed	24h	
<i>Note:</i> Do not use in birds producing eggs for food purposes.					
Gallimycin® PFC	Vétoquinol	Chickens	dw	24h	
<i>Note:</i> Do not use in birds producing eggs for food purposes.					
Gallivac™ HB1 Mass	Merial	Chickens	aer/sp, dw	21d	
IBD Blen®	Merial	Chickens	dw	21d	
IB-Vac-H™	Intervet	Chickens	dw, I.N., I.O.	21d	
Immucox® for Chickens I	Vetech	Chickens	dw, Oral	21d	
Immucox® for Chickens II	Vetech	Chickens	dw, Oral	21d	
Laryngo-Vac™	Wyeth Animal Health	Chickens	I.O.	21d	
Lincomix® 110 Premix	Pfizer	Chickens	In the feed	0d	
<i>Note:</i> Do not use in laying birds.					
Lincomix® Soluble Powder	Pfizer	Chickens	dw	0d	
<i>Note:</i> No drug withdrawal period is required before slaughter of broilers receiving LINCOSOLUBLE powder at the approved concentration (16 mg per litre of drinking water).					
Lincomycin Soluble Powder	Bio Agri Mix	Chickens	dw	0d	
<i>Note:</i> No drug withdrawal period is required before slaughter of broilers receiving LINCOSOLUBLE powder at the recommended concentration (16 mg lincomycin activity per litre of drinking water).					
Lincomycin Spectinomycin 100 Soluble Powder	Bio Agri Mix	Chickens	dw	3d	
<i>Note:</i> Do not use in laying hens.					
Linco-Spectin® 100 Soluble Powder	Pfizer	Chickens	dw	3d	
<i>Note:</i> Do not use in laying birds.					
L-S® 100 Soluble Powder	Pfizer	Chickens	dw	3d	
<i>Note:</i> Do not use in laying birds.					
LT Blen®	Merial	Chickens	dw, I.O.	21d	
LT-Ivax®	Schering-Plough	Chickens	I.O.	21d	
Malathion 50	Vétoquinol	Chickens	Premise	7d	
Marek's Disease Vaccine (HVT)	Merial	Chickens	In ovo, S.C.	21d	
Marek's Disease Vaccine (Rispens CVI 988)	Merial	Chickens	S.C.	21d	
Marek's Disease Vaccine (SB <sub>1</sub> )	Merial	Chickens	In ovo, S.C.	21d	
Marexine™-CA	Intervet	Chickens	S.C.	21d	
Maxiban®	Elanco	Chickens	In the feed	4d	
<i>Note:</i> Do not give to laying hens.					
MBL AE	L.A.H.I.	Chickens	S.C.	42d	
MBL BD3-ND-IB2-REO	L.A.H.I.	Chickens	S.C.	42d	
MBL BD3-REO	L.A.H.I.	Chickens	S.C.	42d	
MBL FC3	L.A.H.I.	Chickens	S.C.	42d	
MBL ND-IB2	L.A.H.I.	Chickens	S.C.	42d	
MBL SE4C	L.A.H.I.	Chickens	S.C.	42d	
MD-Vac® CFL	Wyeth Animal Health	Chickens	S.C.	21d	
Medivit	Vétoquinol	Chickens	dw	5d	
<i>Note:</i> Do not use in laying birds.					
Mildvac-M®	Intervet	Chickens	aer/sp, bd, dw, I.N., I.O.	21d	
M-Ninevax® C	Schering-Plough	Chickens	ww	21d	
Monensin Premix	Bio Agri Mix	Chickens	In the feed		
<i>Note:</i> Do not feed to replacement and laying chickens.					
Monteban® 70	Elanco	Chickens	In the feed		
<i>Note:</i> Do not give to laying hens.					
Neo-Chlor®	Vétoquinol	Chickens	dw	7d	
<i>Note:</i> Do not use in laying birds.					
NeoMed 325	Medprodex	Chickens	dw	7-14d	
Neomix® Soluble Powder	Pfizer	Chickens	dw	7-14d	
Neomycin 325	Vétoquinol	Chickens	dw	7-14d	
Neo Oxymed	Medprodex	Chickens	dw	7d	
<i>Note:</i> Do not feed to laying birds.					
Neotet	Dominion	Chickens	dw	7d	
<i>Note:</i> Do not feed to laying chickens.					

**Chicken Withdrawal Time Chart**
**Withdrawal times listed correspond to label dosages only.**

Product Name	Company Name	Species	Route of Administration	Meat	Milk
Neo-Tetramed <i>Note:</i> Do not use in laying birds.	Medprodex	Chickens	dw	7d	
Neox <i>Note:</i> Do not use in laying birds. Do not use in replacement chickens over 16 weeks of age.	Vétoquinol	Chickens	dw	7d	
New Bronz® <i>Note:</i> Do not administer to laying chickens.	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
New Bronz® MG	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
Newcastle-Bronchitis Vaccine (B1 Strain, Mass. and Conn. Types)	Wyeth Animal Health	Chickens	aer/sp, dw	21d	
Newcastle-Bronchitis Vaccine (LaSota Strain, Mass. Type) LS Mass II™	Wyeth Animal Health	Chickens	dw	21d	
Newcastle-Bronchitis Vaccine (Mass. and Conn. Types)	Merial	Chickens	aer/sp, dw, I.O.	21d	
Newcastle-K <i>Note:</i> Do not administer to laying chickens.	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
Newcastle Vaccine (B1 Strain)	Wyeth Animal Health	Chickens	aer/sp, dw, I.N., I.O.	21d	
Nicarb® <i>Note:</i> Do not feed to laying or breeding hens in production.	Merial	Chickens	In the feed	4d	
Onycin 62.5 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Onycin 250 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Onycin 1000 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Oxy 250 <i>Note:</i> Do not feed to laying birds.	Medprodex	Chickens	dw	7d	
Oxy 1000 <i>Note:</i> Do not use in laying birds.	Jaapharm	Chickens	dw	7d	
Oxysol 62.5 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	7d	
Oxysol-110	Vétoquinol	Chickens	In the feed	7d	
Oxysol-220	Vétoquinol	Chickens	In the feed	7d	
Oxysol 250 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	7d	
Oxysol 440	Vétoquinol	Chickens	In the feed	7d	
Oxysol 1000 <i>Note:</i> This product must not be used in laying poultry.	Vétoquinol	Chickens	dw	7d	
Oxy Tetra-A <i>Note:</i> Eggs taken from treated birds within 60 hours after the latest treatment with this drug must not be used in food.	Dominion	Chickens	dw, In the feed	7d	
Oxytetracycline 50 Granular Premix	Bio Agri Mix	Chickens	In the feed	7d	
Oxytetracycline 100 Granular Premix	Bio Agri Mix	Chickens	In the feed	7d	
Oxytetracycline 200 Granular Premix	Bio Agri Mix	Chickens	In the feed	7d	
Oxy Tetra Forte	Dominion	Chickens	dw, In the feed	7d	
Pabac® <i>Note:</i> Do not vaccinate birds in egg production.	Wyeth Animal Health	Chickens	S.C.	42d	
Paracillin® SP <i>Note:</i> Do not use in laying birds.	Intervet	Chickens	dw	2d	
Poly-tonine A Super Booster No. 1® <i>Note:</i> Eggs obtained from treated birds within 5 days after the latest treatment with this drug must not be used in food.	Vétoquinol	Chickens	dw	5d	
Pro-Gen® 20%	Vétoquinol	Chickens	In the feed	5d	
Pro Vac® 3 ACL™	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
Pro Vac® 4 ACL™	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
PT Blen®	Merial	Chickens	ww	21d	
Quinoxine-S	Vétoquinol	Chickens	dw	4d	
ReoGuard™ L	Merial	Chickens	dw	21d	
Rismavac®	Intervet	Chickens	In ovo, S.C.	21d	
Robenz® Medicated Premix <i>Note:</i> Do not feed to laying hens, replacement and breeding chickens.	Alpharma	Chickens	In the feed	6d	
Rumensin® Premix <i>Note:</i> Do not feed to replacement and laying chickens.	Elanco	Chickens	In the feed		
Sacox® 120 <i>Note:</i> Do not feed to replacement, laying or breeding chickens.	Huvepharma AD	Chickens	In the feed		
Salinomycin 60 Premix <i>Note:</i> Do not feed to replacement, laying or breeding chickens.	Bio Agri Mix	Chickens	In the feed		
Sevin®	Dominion	Chickens	Premise, Topical	7d	
Sevin®	Vétoquinol	Chickens	Topical	7d	
Sodium Sulfamethazine Solution 12.5% <i>Note:</i> This product must not be used in laying poultry.	Dominion	Chickens	dw	12d	

**Withdrawal times listed correspond to label dosages only.**

**Chicken Withdrawal Time Chart**

Product Name	Company Name	Species	Route of Administration	Meat	Milk
Sodium Sulfamethazine Solution 25% <i>Note:</i> This product must not be used in laying poultry.	Dominion	Chickens	dw	12d	
Spectam® Powder <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Spectam® Water Soluble Concentrate <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Stafac® 22 <i>Note:</i> Do not feed to birds producing eggs for human consumption.	Phibro	Chickens	In the feed		
Stafac® 44 <i>Note:</i> Do not feed to birds producing eggs for human consumption.	Phibro	Chickens	In the feed		
Stafac® 500 <i>Note:</i> Do not feed to birds producing eggs for human consumption.	Phibro	Chickens	In the feed		
Sulfa "25" <i>Note:</i> This product must not be used in laying poultry.	P.V.L.	Chickens	dw	12d	
Sulfa 25% Solution <i>Note:</i> Do not use in laying birds.	Bimeda-MTC	Chickens	dw	12d	
Sulfaquinoxaline 19.2% Liquid Concentrate <i>Note:</i> This product must not be used in laying poultry.	Dominion	Chickens	dw	12d	
Sulfaquinoxaline-S	Medprodex	Chickens	dw	4d	
Super Nitro-12 Tablets	Dominion	Chickens	dw	5d	
SVS 510	Merial	Chickens	aer/sp, dw, S.C.	21d	
Terramycin®-50 Premix	Phibro	Chickens	In the feed	7d	
Terramycin®-100 Premix	Phibro	Chickens	In the feed	7d	
Terramycin®-200 Premix	Phibro	Chickens	In the feed	7d	
Tetra 55 <i>Note:</i> Do not use in laying birds.	Jaapharm	Chickens	dw	5d	
Tetra 250 <i>Note:</i> Do not use in laying birds.	Jaapharm	Chickens	dw	5d	
Tetra 1000 <i>Note:</i> Do not use in laying birds.	Jaapharm	Chickens	dw	5d	
Tetracycline 250 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Tetracycline 1000 <i>Note:</i> Do not use in laying birds.	Vétoquinol	Chickens	dw	5d	
Tetracycline Hydrochloride <i>Note:</i> This product must not be used in laying poultry.	Dominion	Chickens	dw	5d	
Tetracycline Hydrochloride <i>Note:</i> Do not administer to laying birds.	P.V.L.	Chickens	dw	5d	
Tetramed 250 <i>Note:</i> Do not feed to laying birds.	Medprodex	Chickens	dw	5d	
Tetramed 1000 <i>Note:</i> Do not feed to laying birds.	Medprodex	Chickens	dw	5d	
Tremor Blen® D <i>Note:</i> Do not vaccinate within 4 weeks of initial egg production.	Merial	Chickens	dw, ww	21d	
Tremvac® <i>Note:</i> Do not vaccinate 35 days prior to onset of lay or during egg production.	Intervet	Chickens	dw	21d	
Tremvac-FP® <i>Note:</i> Do not vaccinate 35 days prior to onset of lay or during egg production.	Intervet	Chickens	ww	21d	
Triplevac®	Intervet	Chickens	aer/sp, dw	21d	
Tri-Reo® <i>Note:</i> Do not administer to laying chickens.	Wyeth Animal Health	Chickens	I.M., S.C.	42d	
Tylan® Soluble <i>Note:</i> Do not use in laying hens.	Elanco	Chickens	dw	1d	
Univax-BD®	Schering-Plough	Chickens	dw, In ovo, S.C.	21d	
Univax®-Plus	Schering-Plough	Chickens	dw, S.C.	21d	
V.A.-Vac® <i>Note:</i> Do not vaccinate birds in egg production.	Wyeth Animal Health	Chickens	S.C.	21d	
Vibiomed Booster <i>Note:</i> Eggs obtained from treated birds within 5 days after the latest treatment with this drug must not be used in food.	Medprodex	Chickens	dw, In the feed	5d	
Vineland® BD3-ND-IB1-REO	L.A.H.I.	Chickens	I.M., S.C.	42d	
Vineland® BD4-REO	L.A.H.I.	Chickens	I.M., S.C.	42d	
Vineland® Chick Poly Banco™	L.A.H.I.	Chickens	dw, I.N., I.O.	21d	
Vineland® FC-3	L.A.H.I.	Chickens	S.C.	42d	
Vineland® HVT+Rispens	L.A.H.I.	Chickens	S.C.	21d	
Vineland® HVT+SB1	L.A.H.I.	Chickens	In ovo, S.C.	21d	
Vineland® MLT	L.A.H.I.	Chickens	aer/sp, I.N., I.O.	21d	
Vineland® ND-IB1	L.A.H.I.	Chickens	I.M., S.C.	42d	

**Chicken Withdrawal Time Chart****Withdrawal times listed correspond to label dosages only.**

<b>Product Name</b>	<b>Company Name</b>	<b>Species</b>	<b>Route of Administration</b>	<b>Meat</b>	<b>Milk</b>
Vineland® PP	L.A.H.I.	Chickens	ww	21d	
Vineland® Vi Bursa-G™	L.A.H.I.	Chickens	dw	21d	
Vineland® Vi So Bronc®	L.A.H.I.	Chickens	aer/sp, dw, I.N., I.O.	21d	
Vineland® Vi-Tremox™	L.A.H.I.	Chickens	ww	21d	
Virginiamycin 44 Premix	Bio Agri Mix	Chickens	In the feed		
<b>Note:</b> Do not use in birds producing eggs for food purposes.					
Zoamix®	Huvepharma AD	Chickens	In the feed		
<b>Note:</b> Do not administer to laying chickens. Do not feed to replacement chickens after the first 14 weeks of life.					